



JIRÍ MLÍKOVSKÝ

## METHODOLOGICAL PROBLEMS OF THE CONTEMPORARY SYSTEMATICS OF THE FAMILY HOMINIDAE

**ABSTRACT.** — *Methodological problems of the contemporary hominid systematics are pointed out and briefly discussed from the point of view of general systematics. The necessity is stressed (1) to distinguish between the evolution (of organisms) and phylogeny (of taxa), (2) to distinguish between the phylogenetic and identification diagnoses, and (3) that taxonomic decisions should be made within properly arranged groups only, i.e. here within the Hominoidea, not within the Hominidae.*

**KEY WORDS:** *Systematics — Methodological problems — Hominidae.*

Besides evolutionary morphological and functionally morphological studies, the contemporary anthropology pays a great deal of attention to systematics. In spite of the relatively great scientific potential (as compared e.g. with zoology) devoted to this research, our knowledge is still surprisingly minute. When trying to analyze the causes of this situation we can see that hominid systematics was completely isolated from the zoological one for a long time and has developed its methods independently. The attempts of zoologists to surpass this state were viewed negatively by the anthropologists, and it resulted in a sceptical opinion of zoologists on the anthropological systematists and their efforts. There were even suggestions to exclude hominids from the field of zoosystematics (e.g. Simpson, 1945: 188).

In the last years this situation has markedly changed to the better. The anthroposystematists have begun to be interested in the methods of zoological systematics and even in the methodology of general systematics, and, on the other hand, the zoologists (especially primatologists) have increasingly begun to deal with hominids which ceased to be a strict taboo for them, though a mistrust towards zoologists and their methods from the anthropologists side still continues.

The exact reasons for this change of opinion could be explained only by a detailed historic analysis (I am sure a very useful one). The chief factors of this change were probably on the one hand the general trend of the contemporary biology to synthesize all available information (see e.g. Čepi-

kov, 1975, Novák, 1979, Fedorov, 1979), and on the other the crisis of the anthroposystematics which forced this scientific field to join that general process. Due to this development, a mixture of different and often contradictory opinions and trends occurs in the contemporary anthroposystematics which the non-specialist can hardly grasp. I tried therefore to analyze this complicated situation from the viewpoint of the general systematics.

The analysis of methodology of the contemporary systematics discovered a number of methodological shortcomings responsible for the complexity of this situation.

Apparently the basic difficulty is the inconsequent distinguishing among the problems of phylogeny, evolution, and identification. We should keep in mind that not all of the methods of study of the evolution of organisms (or their characters) may be used for the study of the phylogeny, i.e. the historical development of taxa (or their characters), and vice versa. Unfortunately, a detailed discussion of this problem which is necessary for any exact understanding of the given differences is not possible here because it would exceed the limits of the present paper and because the solution of this problem has not been reached even from the point of view of the general systematics till now. On the other hand, the identification deals with the determination of individual finds using diagnoses of taxa for these purposes. However, these identification diagnoses considerably differ from the phylogenetic ones by their significance and content. Unfortunately, neither do the zoological and botanical

systematists realize this difference for the present. The phylogenetic diagnosis should define the phylogenetic position of the given taxon while the identification diagnosis is concerned with the identification of individual finds.

The problem of the diagnosis of taxa is connected with the problem of nomenclature which is common for both the anthroposystematics and the systematics of other groups of organisms. Systematists conceive the taxa as classical sets (cf. e.g. Gregg 1954, Buck and Hull 1966) the denotation of which is sufficiently elaborated, but which are inconsistent with the taxonomic continuum which in turn is characteristic for the phylogeny. A number of investigators attempted to solve this inconsistency without breaking the paradigm of transformation of the taxonomic continuum into the classical sets. Various systems were proposed, as e.g. the overlapping taxa (Michener 1963), bigeneric names (Tobias 1969, 1973), successional species (Imbrie 1957), sister groups (Hennig 1950, 1966), etc. A new approach to the problem of taxonomic continuum represents only the punctuated equilibrium theory (Eldredge and Gould 1972, Gould and Eldredge 1977). As compared with the classical gradualistic theory this theory brings some significant improvements, though some details are questionable. Together with the fact that taxa represent fuzzy sets, not classical ones (see Mlíkovský 1979, 1983, in prep.), this theory suggests an escape from the discussed dilemma. However, new problems arise because direct application of the systematic nomenclature to the fuzzy sets is not possible. Consequently a new method of biological denotation of taxa conceived as fuzzy sets will have to be elaborated. Although this problem has not yet been solved, some analyses indicate (Mlíkovský in prep.) that such a solution is possible and that the currently used nomenclature will need not to be substantially changed. In connection with these deliberations the necessity of a strict adherence to the International Rules of Zoological Nomenclature should be stressed; there are anthropologists who often do not observe these Rules consequently.

The next general problem of the systematics is the problem of categories, i.e. the problem of an ascription of taxa to categories. This problem is especially acute in the anthropology due to the discussion as to which organism can be and which cannot be denoted as the man, i.e. from the paleo-anthropological point of view, which fossil taxa can be and which cannot be classified into the genus *Homo*. This discussion, however, deals with a pseudoproblem because there is no sense in ascribing objects to the sets (i.e. taxa here) in the case when these sets are not explicitly defined, and that are the genera of hominids not. The anthropological taxonomy should therefore first solve the problem of phylogenetic definitions of its taxa, then introduce identification diagnoses on the basis of these definitions, and finally it may ascribe individual finds or phena to the individual taxa on the basis of these identification diagnoses. Yet from the point of view of the general systematics there should be

the highest possible equivalence of the definition of the anthropological categories and taxa with the definitions of these units in other biological disciplines (Mlíkovský 1979, in prep.).

The last methodological problem of the general systematics which shall be discussed here and which leads to the problems of special systematic methods is the relation between similarity and affinity. This problem is based on the fact that an observable quality is similarity while relationships (affinity) could be assessed only indirectly, on the basis of similarity and the analysis of its causes (Mlíkovský 1979, in prep.). All attempts to solve these problems could be divided into 3 groups: 1. the classical ("evolutionary") systematics, 2. the cladistic ("phylogenetic") systematics, and 3. the numerical ("phenetic") systematics.

Classical systematics (Plate 1914, Rensch 1934, Schilder 1952, Mayr et al. 1953, Simpson 1961, Blackwelder 1967, Mayr 1969, Smirnov 1969, Bărănescu 1973, Ross 1974, Zarenkov 1976, Mertens and Lines 1978) is based on a subjective decision as whether the similarity of two taxa expresses their relationships or whether it is brought about by convergence. Many rules were introduced later in order to make the decision-making process more objective (see the literature cited above, and further especially Remane 1956). This trend, however, resulted rather in a standardization than in the objectivization of the process.

A breakthrough in this convention was attempted only by Hennig (1950, 1965, 1966) with his cladistic systematics (see also Schlee 1971, Dupuis 1979, and Wiley 1981), in which he supposes that there exists a correlation between the evolutionary splitting of the given line and the time of the scientific fixation of groups which arose from it, and that the relationships can be derived directly from the evolution of individual characters. Although this method is often used in zoology (but almost unknown in botany — see Bremer and Wanntorp 1978) for many reasons it cannot be accepted (cf. e.g. Inger 1967, Mayr 1969, 1974, Van Valen 1978, Eichler 1978, Thenius 1979). Fortunately it is used very infrequently in anthroposystematics (e.g. Eldredge and Tattersall 1975, Delson 1978, Schwartz et al. 1978). However, interesting results could be provided by the so-called character compatibility analysis (Estabrook 1972, McMorris 1975, Estabrook et al. 1975, 1976a, b, 1977) which was derived from the cladistic analysis using numerical methods. However, this method has not been used either in the anthropology or in the primatology, yet.

The third group of the systematic methods is the so-called numerical systematics (Sokal and Sneath 1963, Jardine and Sibson 1971, Sneath and Sokal 1973, Leuschner 1974, Clifford and Stephenson 1975). It is based on the presumption that the general similarity is directly correlated with the relationships (provided that a sufficient number of characters was analyzed). It was, however, experimentally shown that this presumption is incorrect (e.g. Boyce 1964, Eades 1965, Minkoff 1965, Mayr

1965, 1969, Sokal and Michener 1967, Kendrick and Weserup 1967, Crovello 1968, 1969, Rohlf 1972). Despite this evidence the numerical systematics is very popular in biology, perhaps owing to its seeming exactness and scientific value. In anthropology this method is dealt with by, e.g., Oxnard (1968), Bilsborough (1971), Kowalski (1972), Corruccini et al. (1976, 1979), Corruccini (1978), and Corruccini and Henderson (1978).

Consequently, none of these systematic methods is fully satisfactory, and this fact should always be kept in mind during the interpretation of results. Some significant improvements should be introduced by a new systematic method denoted as logical systematics (Mlíkovský 1979) which I am elaborating at present.

But none of the systematic methods could yield correct results unless the two following conditions are met: 1. the proper delimitation of phena, and especially 2. the proper arrangement of the group, in which phylogeny is studied. Every analyzed group should be monophyletic and its elements should be real. In the case of hominids I presume, e.g., that it is not entirely correct to analyze the phylogeny or the evolution within the framework of the family Hominidae because this family is not clearly separated from other families of the superfamily Hominoidea. Consequently there is a possibility that some taxa and/or phena included in this family at present are not members of it, whereas some other taxa (phena) which have been included in another family of the superfamily Hominoidea belong to this group, as it may be shown in the discussion about the genera *Gigantopithecus* (Frayer 1973, Eckhard 1975, Simons 1978), and *Ramapithecus* (Conroy 1972, Vogel 1975, Aguirre 1975, Conroy and Pilbeam 1975, Simons 1976, 1978, Frayer 1978, Greenfield 1979). On the other hand the superfamily Hominoidea is separated from other superfamilies of primates clearly enough so that the possibility of the complications mentioned above is not too high and the systematic studies within this group could therefore be made with a hope to some purpose.

At the end I would like to stress the necessity of a synthetic approach to the solution of the anthroposystematic problems. For a successful development of the anthroposystematics it is necessary to interconnect organically its paleontological and neontological parts (as happens now in zoology) and to remove the barrier between the anthroposystematics and the systematics of other organisms, because despite every plant and animal group (i.e. consequently also the hominids) has its own concrete specific features, the principles of the general systematics hold for all of them.

#### ACKNOWLEDGMENTS

I would like to thank my colleagues D. Zemková, V. Přívratský, and especially V. Vančata for introducing me into the problems of anthropology, for providing me with anthroposystematic literature

and for discussion of this paper, for which I am greatly indebted also to my colleague K. Zemek.

#### REFERENCES

- AGUIRRE E., 1975: *Kenyapithecus* and *Ramapithecus*. In: *Paleoanthropology, morphology and paleoecology*. Ed. R. H. Tuttle, Pp. 99–103. The Hague, Mouton.
- BARANESCU P., 1973: *Principiile și metodele zoologiei sistematice*. Acad. Republ. Soc. România, Bucuresti. 219 pp.
- BILSBOROUGH A., 1971: Evolutionary change in the hominid maxilla, *Man*, 6: 473–486.
- BLACKWELDER R. E., 1967: *Taxonomy*. Wiley, New York. 698 pp.
- BOYCE A. J., 1964: The value of some methods of numerical taxonomy with reference to hominoid classification. In: *Phenetic and phylogenetic classification*. Ed. V. H. Heywood and J. McNeil. Pp. 45–65.
- BREMER K., WANNTORP H.-E., 1978: Phylogenetic systematics in botany. *Taxon*, 27: 317–329.
- BUCK R. C., HULL D. L., 1966: The logical structure of the Linnean hierarchy. *Syst. Zool.*, 15: 97–111.
- ČEPIKOV M. G., 1975: *Integracija nauki*. Moskva: Mysl'. 244 pp.
- CLIFFORD H. T., STEPHENSON W., 1975: *An introduction to numerical classification*. Academic Press, New York. 229 pp.
- CONROY G. C., 1972: Problems in the interpretation of *Ramapithecus*: With special reference to anterior teeth reduction. *Amer. J. phys. Anthropol.*, 37: 41–48.
- CONROY G. C., PILBEAM D. R., 1975: *Ramapithecus*: A review of its hominid status. In: *Paleoanthropology, morphology, and paleoecology*. Ed. R. H. Tuttle. Pp. 59–86. The Hague, Mouton.
- CORRUCCINI R. S., 1978: Morphometric analysis: Uses and abuses. *Yearbook phys. Anthropol.*, 21: 134–150.
- CORRUCCINI R. S., CIOCHON R. L., HENRY H. M., 1976: The postcranium of Miocene hominoids: Were dryopithecines merely "dental apes"? *Primates*, 17: 205–223.
- CORRUCCINI R. S., CRONIN J. E., CIOCHON R. L., 1975: Scaling analysis and congruence among anthropoid primate macromolecules. *Human Biol.*, 51: 167–185.
- CORRUCCINI R. S., HENDERSON A. M., 1978: Multivariate dental allometry in primates. *Amer. J. phys. Anthropol.*, 48: 203–208.
- CROVELLO T. J., 1968: The effect of change of number of OTUs in a numerical taxonomic study. *Brittonia*, 20.
- CROVELLO T. J., 1969: Effects of change of characters and of number of characters in numerical taxonomy. *Amer. Midl. Nat.*, 81: 68–86.
- DELSON E., 1978: Models of early hominid phylogeny. In: *Early hominids of Africa*. Ed. C. Jolly. Pp. 517–541. London, Duckworth.
- DUPUIS C., 1979: La "systematique phylogénétique" de W. Hennig (historique, discussion, choix de references). *Cah. nat.*, 34: 1–69.
- EADES D. C., 1965: The inappropriateness of the correlation coefficient as a measure of taxonomic resemblance. *Syst. Zool.*, 14: 98–100.
- ECKHARD R. B., 1975: *Gigantopithecus* as a hominid. In: *Paleoanthropology, morphology, and paleoecology*. Ed. R. H. Tuttle. Pp. 105–129. The Hague: Mouton.
- EICHLER W., 1978: Kritische Einwände gegen die Hennig'sche kladistische Systematik. *Biol. Rdsch.*, 16: 175–183.
- ELDRIDGE N., GOULD S. J. 1972: Punctuated equilibria: An alternative to phyletic gradualism. In: *Models in paleobiology*. Ed. T. J. M. Schopf, Pp. 82–115. San Francisco, Freeman, Cooper, and Co.
- ELDRIDGE N., TATTERSALL I., 1975: Evolutionary models, phylogenetic reconstructions and another look at hominid phylogeny. *Contrib. Primatol.*, 5: 218–242.
- ESTABROOK G. F., 1972: Cladistic methodology: A discussion of the theoretical basis for the induction of evolutionary history. *Annu. Rev. Ecol. Syst.*, 3: 427 to 456.

- ESTABROOK G. F., JOHNSON C. S., Jr., MORRIS F. R., 1975: An idealized concept of the true cladistic character. *Math. Biosci.*, 23: 263–272.
- ESTABROOK G. F., JOHNSON C. S., Jr., MORRIS F. R., 1976a: An algebraic analysis of cladistic characters. *Discrete Math.* 16: 141–147.
- ESTABROOK G. F., JOHNSON C. S., Jr., MORRIS F. R., 1976b: A mathematical foundation for the analysis of cladistic character compatibility. *Math. Biosci.*, 29: 181–187.
- ESTABROOK G. F., STRAUCH J. G., Jr., FIALA K. L., 1977: An application of compatibility analysis to the Blackith's data on orthopteroid insects. *Syst. Zool.*, 26: 269–276.
- FEDOROV V. M., 1979: *Sintetičeskije tendencii v sovremenom estestvoznanii*. Moskovskij gosudarstvennyj universitet, Moskva. 117 pp.
- FRAYER D. W., 1973: Gigantopithecus and its relationships to Australopithecus. *Amer. J. phys. Anthropol.*, 39: 413–426.
- FRAYER D. W., 1978: The taxonomic status of Ramapithecus. In: *Krapinski pračovjek i evolucija hominida*. Ed. M. Malez. Pp. 255–268.
- GOULD S. J., ELDRIDGE N., 1977: Punctuated equilibria: The tempo and mode of evolution reconsidered. *Paleobiology*, 3: 115–151.
- GREENFIELD L. O., 1979: Adaptive patterns of Ramapithecus. *Amer. J. phys. Anthropol.*, 50: 527–548.
- GREGG J. R., 1954: *The language of taxonomy. An application of symbolic logic to the study of classificatory systems*. Columbia University Press, New York. 70 pp.
- HENNIG W., 1950: *Grundzüge einer Theorie der phylogenetischen Systematik*. Deutscher Zentralverlag, Berlin. 370 pp.
- HENNIG W., 1965: Phylogenetic systematics. *Annu Rev. Entomol.*, 10: 96–116.
- HENNIG W., 1966: *Phylogenetic systematics*. University of Illinois Press, Urbana. 263 pp (reedition: ibidem 1979).
- IMBRIE J., 1957: The species problem with fossil animals. In: *The species problem*. Ed. E. Mayr. Pp. 125–153. New York, American Association for the Advancement of Science.
- INGER R. F., 1967: The development of a phylogeny of frogs. *Evolution*, 21: 369–384.
- JARDINE N., SIBSON R., 1971: *Mathematical taxonomy*. Wiley, New York. 286 pp.
- KENDRICK W. B., WESERUP L. K., 1966: Attempting neo-adansonian computer taxonomy at the ordinal level in the Basidiomycetes. *Syst. Zool.*, 15: 307–329.
- KOWALSKI C., 1972: A commentary on the use of multivariate statistical methods in anthropometrical research. *Amer. J. phys. Anthropol.*, 36: 119–132.
- LEUSCHNER D., 1974: *Einführung in die numerische Taxonomie*. Fischer, Jena.
- MAYR E., 1965: Numerical phenetics and taxonomic theory. *Syst. Zool.*, 14: 73–97.
- MAYR E., 1969: *Principles of systematic zoology*. McGraw-Hill, New York. 428 pp.
- MAYR E., 1974: Cladistic analysis or cladistic classification. *Z. zool. Syst. Evol.forsch.*, 12: 95–128.
- MAYR E., LINSLEY E. G., USINGER R. L., 1953: *Methods and principles of systematic zoology*. McGraw-Hill, New York. 336 pp.
- McMORRIS F. R., 1975: Compatibility criteria for cladistics and qualitative taxonomic characters. *Proc. int. Conf. numer. Taxonomy* 8: 399–415.
- MERTENS T. R., LINES J. L., 1978: *Principles of biosystematics*. 2nd ed. Illustr. Educational Methods, Chicago. 142 pp.
- MICHENER C. D., 1963: Some future developments in taxonomy. *Syst. Zool.*, 12: 151–172.
- MINKOFF E. C., 1965: The effects on classification of slight alternations in numerical technique. *Syst. Zool.*, 14: 196–213.
- MLÍKOVSKÝ J., 1979: *Obecné problémy evoluční systematiky*. Unpublished thesis, Charles University, Praha.
- MLÍKOVSKÝ J., 1983: Zur Methodologie phylogenetischer Rekonstruktionen in der Paläontologie. *Schriegenr. geol. Wiss.* 19/20: in press.
- NOVAK V. J. A. 1979: Vývoji vědy jako formy společenského vědomí a přirozená soustava věd. *Filoz. Čas.*, 27: 824–836.
- OXNARD C. E., 1968: Primate evolution. A method of investigation. *Amer. J. phys. Anthropol.*, 28: 289–302.
- PLATE L., 1914: Prinzipien der Systematik mit besonderer Berücksichtigung des Systems der Tiere. In: *Kultur der Gegenwart III (4/4)*. Ed. P. Hinneberg. Pp. 92 bis 164.
- REMANE A., 1956: *Die Grundlagen des natürlichen Systems der vergleichenden Anatomie und Systematik*. Akademische Verlagsgesellschaft, Leipzig. 400 pp.
- RENSCH B., 1934: *Kurze Anweisung für zoologisch-systematische Studien*. Akademischer Verlag, Leipzig. 116 pp.
- ROHLF F. J., 1972: An empirical comparison of three ordination techniques in numerical taxonomy. *Syst. Zool.*, 21: 271–280.
- ROSS H. H., 1974: *Biological systematics*. Addison-Wesley, Reading. 345 pp.
- SCHILDER F. A., 1952: *Einführung in die Biotaxonomie*. Fischer, Jena. 162 pp.
- SCHLEE D., 1971: Die Rekonstruktionen der Phylogenese mit Hennigs Prinzip. *Aufs. Red. senckenberg. naturforsch. Ges.*, 20: 1–62.
- SIMONS E. L., 1976: Relationships between Dryopithecus, Sivapithecus and Ramapithecus and their bearing on hominid origins. In: *Les plus anciens hominidés*. Ed. P. V. Tobias and Y. Coppens. Paris, C.N.R.S. Pp. 36 to 59.
- SIMONS E. L., 1978: Diversity among the early hominids: A vertebrate paleontologist's viewpoint. In: *Early hominids of Africa*. Ed. C. J. Jolly, London: Duckworth. Pp. 543–566.
- SIMPSON G. G., 1945: The principles of classification and a classification of mammals. *Bull. amer. Mus. nat. Hist.*, 85: 3–350.
- SIMPSON G. G., 1961: *Principles of animal taxonomy*. Columbia University Press, New York, 247 pp.
- SMIRNOV E. S., 1969: Taksonomičeskij analiz. Moskovskij gosudarstvennyj universitet, Moskva. 187 pp.
- SNEATH P. H. A., SOKAL R. R., 1973: *Numerical taxonomy. The principles and practice of numerical classification*. Freeman, San Francisco, 573 pp.
- SOKAL R. R., MICHENER C. D., 1967: The effect of different numerical techniques on the phenetic classification of bees of the Hoplitis complex (Megachilidae). *Proc. linn. Soc. London*, 178: 59–74.
- SOKAL R. R., SNEATH P. H. A., 1963: *Principles of numerical taxonomy*. Freeman, San Francisco. 359 pp.
- THENIUS E. 1979: Hennigs phylogenetische Systematik und paläontologische Befunde. *N. Jb. Geol. Paläontol.*, 1979 (9): 406–414.
- TOBIAS P. V., 1969: Bigeneric nomina: A proposal for modification of the rules of nomenclature. *Amer. J. phys. Anthropol.*, 31: 103–106.
- TOBIAS P. V., 1973: Darwin's prediction and the African emergence of the genus Homo. In: *L'origin dell'Uomo*. Pp. 63–85.
- VAN VALEN L., 1978: Why not to be a cladist. *Evol. Theory*, 3: 285–299.
- VOGEL C., 1975: Remarks on the reconstruction of the dental arcade of Ramapithecus. In: *Paleoanthropology, morphology, and paleoecology*. Ed. R. H. Tuttle. The Hague: Mouton. Pp. 87–98.
- WILEY E. O., 1981: *Phylogenetics. The theory and practice of phylogenetic systematics*. Wiley and Sons, New York. 439 pp.
- ZARENKOV N. A., 1976: *Lekcii po teorii sistematiki*. Izdatelstvo Moskovskogo Universiteta, Moskva. 140 pp.

J. Mlíkovský  
Dept. of Evolutionary Biology  
Institute of Microbiology  
Czechoslovak Academy of Sciences  
Na Folimance 5  
CS – 120 00 Praha 2  
Czechoslovakia